

STUDY ON RELATIONSHIP BETWEEN HORMONES AND AGGRESSION IN DIFFERENT ANIMALS AND HUMAN

Suresh Bhardwaj¹ Dr. D.B. Singh²

¹Research Scholar, CMJ University, Shillong, Meghalaya

²Deptt. Of Management, Lord Krishna College Of Engineering, Ghaziabad (U.P.)

ABSTRACT

To compare brain activity in 41 convicted violent offenders to activity in 41 age matched control subjects. They found that the people convicted of murder had reduced activity in the prefrontal cortex and increased activity in sub-cortical regions such as the thalamus. This finding fits nicely with previous research showing that the damage to the prefrontal cortex impairs decision making and increasing impulsive behavior. Indeed, Raine's work is perhaps the best evidence yet that impaired brain functioning may underlie some types of violent aggression. However, it is important to remember that his subjects lie at the extreme end of a spectrum and may not be typical of most aggressors.

Key words: *prefrontal cortex, evidence, aggression*

INTRODUCTION

Although it seems that everybody understands what aggression is, yet there is considerable disagreement about how to define it. The published literature on human aggression show variations in definitions proposed. From the review of literature related to the conceptualization of human aggression, it is clear that three important distinctions should be made for proper understanding of the concept. The first is whether it should be defined simply in terms of hurtful behaviour, or a person's hurtful intentions should be taken into consideration. Behaviourists have taken aggression as any behaviour that hurts others. The advantage of this definition is that the behaviour itself determines whether or not an act is aggressive. But this definition ignores the intention of person who does this act. Intentions have a central role in the judgments about aggression. In this context aggression can be taken as any action that is intended to hurt others.

REVIEW OF LITERATURE

Testosterone is another attractive candidate for mediating aggression because males in of all ages, races and cultures are more physically aggressive than their female counterparts. In animals, testosterone is linked to social aggression. Reducing testosterone in the alpha male by castrating him eliminates his dominant social status, and restoring testosterone through injection causes him to regain his social status. However, administering testosterone to males with less social status does not usually allow them to take over the alpha male position, indicating that there is not a direct relationship between testosterone and position in the dominance hierarchy. There is some evidence in humans that high testosterone males are more likely to be socially aggressive, but no evidence that they are necessarily more violent. Often they are successful in professions that thrive on competition, such as successful leading of a company, running for president, or pursuing a sports career. Also, a few psychologists have suggested that females are not necessarily less aggressive than males; rather, they display a different kind of aggression. Females are more likely to show non-violent types of aggression such as ostracizing their peers or spreading false rumors with the intent to cause pain. Thus, while there does seem to be a connection between testosterone and physical aggression, a person's testosterone level will not necessarily be a good predictor of aggressive behavior.

MATERIAL AND METHOD

Measures of brain functioning such as the EEG have long suggested that violent criminals have impaired neurological processes, but the recent advancement of neuro imaging techniques has allowed researchers to examine violent offenders' brains in more detail. Adrian Raine and colleagues have conducted the largest and most thorough study to date, in which they used positron emission tomography (commonly called a PET scan) to compare brain activity in 41 convicted violent offenders to activity in 41 age matched control subjects. They found that the people convicted of murder had reduced activity in the prefrontal cortex and increased activity in sub-cortical regions such as the thalamus. This finding fits nicely with previous research showing that the damage to the prefrontal cortex impairs decision making and increasing impulsive behavior. Indeed, Raine's work is perhaps the best evidence yet that impaired brain functioning may underlie some types of violent aggression. However, it is important to remember that his subjects lie at the extreme end of a spectrum and may not be typical of most aggressors. Also,

there are plenty of examples of people with prefrontal cortex damage who do not commit violent acts, so PET scans cannot be used to ferret out potential murders.

Table 1 Inter-correlation Matrix among Females

	A	B	C	D	E	F	G	H	I	J	O	Q2	Q3	Q4	Ex	Inh	Mob	Va	Pa	Ata
A	-	.0	.0	.1	.4	.9	.0	.5	.6	.3	.4	-.1	.04	.07	.11	**	.07	.08	.11	.22*
B			.0	.0	.4	.5	.2	.1	.6	.5	.7	.08	.06	.16	.06	.13	.05	.01	0	.08
C				.3	.4	.0	.2	.0	.0	.1	.8	*	.26	.40	.26	.29	.09	.20	-.25*	.22*
D					.0	.8	.4	.0	.1	.0	.6	-.02	.44	.47	.26	.35	.14	.15	.36*	.32**

E									.01	.25*	.07	.05*	.03	.15	.10	.11	.05	.09	.11	.02	0
F										.13*	.03	.03	.02	.41**	.09	.03	.25*	.02	0	.06	.06
G										.08	.05	.09	.07	.35**	.25*	.13	.32**	.13	.05	.26*	.21*
H										.09	.05	.07	.09	.25*	.06	.33**	.02	.01	.07	.03	.02
I										.01	.04	.07	.07	.11	.03	.11	.12	-.05	.02	-.04	.02
J										.08	.45**	.07	.08	.05	.09	.06	.07	.09	.12	-.1	.11
O											.14	.14	.14	.26**	.28**	.43**	.15	.03	.05	.15	.16
Q2														.12	.12	-	.05	.21	-	-.07	.04

											.11	*	.04			
Q3											- .36**	.42**	.40**		- .03	.20*
Q4											- .40**	-.39**	- .15		.34*	.22*
Ex												.37**	.21*	- .09	.29**	
Inh													.25*	.28**	.38*	.25*
M ob														- .06	.26*	-.15
Va															.19	.09
Pa																.42**
At a																

CONCLUSION

Several lines of converging evidence indicate that the neurotransmitter serotonin plays a key role in mediating aggressive and violent behavior. Mice with a selective knockout of the serotonin 1B receptor show an increase in aggression. Similarly, depleting serotonin levels in vervet monkeys increases their aggressive behavior, whereas augmenting serotonin levels reduces aggression and increases peaceable interactions like grooming. Serotonin has also been implicated in human aggression. For example, pharmacological interventions that augment serotonergic efficacy have been shown to reduce hostile sentiment and violent outbursts in aggressive psychiatric patients. Also, people with a history of impulsively violent behavior, such as arsonists, violent criminals, and people who die by violent methods of suicide show low levels of the serotonin in their cerebral spinal fluid. These findings represent an interesting correlation, but it is important to remember that the direction of effect is unclear. It may be that aggressive behavior induces low serotonin levels in the cerebral spinal fluid rather than vice versa.

REFERENCES

- Bushman, B.J. development(1996). Individual differences in the extent and of aggressive cognitive-associative networks. Personality and Social Psychology Bulletin, 22, 811-819.*
- DeRidder, R., Schruijer, S.G.L., & Tripathi, R.C. (1992). Norm violation as a precipitating factor of negative intergroup relations. In R. DeRidder & R.C. Tripathi (Eds.), Norm violation and intergroups relations (pp. 3-37).*
- Karma, Akko P.; Visser, Lieuwe & Peeters, Allerd. (Utrecht U, Inst of social & Organizational Psychology, Netherlands). Sociable and aggressive dominance: Personality differences in leadership style? Leadership Quarterly, 1993(Spr), Vol. 4(1), 45-64.*
- Ohannessian, Christine McCauley & Hesselbrock, Victor M.(U Connecticut Heath Ctr, Dept of Psychiatry, Farmington) Temperament and personality typologies in adult off spring of alcoholics. Journal of Studies on Alcohol, 1995(May), vol. 156(3), 318-327.*
- Parker, J.G., & Asher, S.R. (1987). Peer relations and later personal adjustment: Are low- accepted children at risk? Psychological Bulletin, 102, 357-389.*
- Paterson, Gill & Sanson, Ann. (Australian Catholic U. Dept of Pederson, N.L., Lichtenstein, P., Plomin, R., DeFaire, U., Mc-Clearn, G.E., & Matthews, K .A. (1989). Genetic and environmental influences for Type A-like measures and related trails: A study of twins reared apart and twins reared together, Psychosomatic Medicine, 51, 428-440.*

Rothbart, Mary K.; Ahadi, Stephan A. & Hershey, Karen L. (U Oregon, Eugene) Temperament and social behavior in childhood.

Rule, B.G., Taylor, B.R., & Dobbs, A.r (1987). Priming effects of heat on aggressive thoughts. Social Cognition, 5, 131-143.

Ruston, J.P., Fulker, D.W., Neale, M.C., Nias, D.K.13., & Eysenck, H.J. (1986), Altruism and aggression: The heritability of individual differences. Journal of personality and social psychology, 50, 1192-1198.

Rutter, Michael. (MRC, Child Psychiatry Unit, England) Temperament, Personality, and personality disorder. British Journal of Psychiatry, 1987 (Apr). vol. 150, 443-458.

Sakic, Vlado & Fulgosi-Masnjak, Rea. (Migration Research Ctr, Smith, T. W. (1992). Hostility and health: Current status of psychosomatic hypothesis. Health Psychology, 11, 139-150.

Spielberger, C.D., Johnson, E.H., Russell, S.F., Crane, 12..1., Jacobs, G..i., & Worden, T.,I. (1985). The experience and expression of anger: Construction and validation of an anger expression scale. In M. A. Chesney & R.H. Ronsenman. (Eds.), Anger and hostility irr cardiovascular and behavioral disorders (pp. 5-30). Washington, DC: Hemisphere.